




\*LAP free : Land Animal Protein free

# B-NATURE GROWER

Extruded pellet for ongrowing **Trout & Others Salmonids**



- Feed **usable in Organic salmonids farming** conditions
- **High level of digestible energy, allowing flexibility** and year-round use
- **High DP/DE ratio**
- **Ensures a good nutritional balance** throughout the cycle and **preserves flesh quality**
- Range available with or without pigments

B-NATURE GROWER				
Commerciale Ref.	4	5	7	9
Diameter (mm)	4	5,5	7,5	9,5
Presentation	Pellet			
Live weight (g)	50 to 100 g	100 to 400	400 to 1 800	1 800 to 2 500

This fish feed can be used in **Organic farming**, in accordance with the regulation (CE) n°834/2007, (CE) n°889/2008 et (CE) 710/2009. This approach is controlled by FR-BIO 10.

### PACKING

15 kg, 20 kg, 25 kg bag, big bag or bulk

*Store feed in a cool and dry place*

### FLOATING TYPE

Sinking, half-floating and floating

### INDICATIVE NUTRITIONAL PROFILE

		B-NATURE GROWER
<b>Protein</b>	(%)	<b>45</b>
<b>Fats</b>	(%)	<b>17</b>
<b>Digestible Energy</b>	(MJ/Kg)	<b>18,2</b>
<b>DP / DE</b>	(g/MJ)	22,7
<b>Gross Energy</b>	(MJ/Kg)	20,8
<b>Fibre</b>	(%)	1,5
<b>Ash</b>	(%)	13,2
<b>Phosphorus</b>	(%)	1,7

### VITAMINS INCORPORATION

Vit. A (UI/Kg)	10 000
Vit. D3 (UI/Kg)	1 750
Vit. E (mg/Kg)	200
Vit. C (mg/Kg)	250

### SUSTAINABILITY

Here at Le Gouessant, we believe that aquaculture is sustainable if performance, quality and preservation of marine resources are combined, that is why we are Global GAP certified.



## FEEDING TABLE

Live Weight (g)		Feed	P	F	Diameter		Feeding rate (% of Biomass / day) following water temperature																					
From	To				N°	(mm)	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20					
50	100	<b>B-NATURE GROWER</b>	<b>45</b>	<b>17</b>	<b>4</b>	<b>4</b>	1,10	1,18	1,26	1,36	1,44	1,54	1,62	1,71	1,81	1,88	1,94	1,97	1,98	1,96	1,92	1,79	1,62					
100	150				<b>5</b>	<b>5,5</b>	0,97	1,05	1,13	1,22	1,31	1,40	1,49	1,58	1,66	1,73	1,79	1,84	1,86	1,83	1,77	1,64	1,49					
150	200				<b>5</b>	<b>5,5</b>	0,89	0,96	1,05	1,13	1,21	1,29	1,37	1,47	1,55	1,64	1,70	1,75	1,76	1,73	1,65	1,53	1,37					
200	400				<b>5</b>	<b>5,5</b>	0,85	0,92	0,99	1,08	1,15	1,22	1,30	1,37	1,46	1,54	1,61	1,64	1,65	1,62	1,54	1,43	1,30					
400	500				<b>7</b>	<b>7,5</b>	0,77	0,84	0,91	0,99	1,06	1,13	1,21	1,27	1,35	1,42	1,48	1,51	1,51	1,48	1,42	1,32	1,21					
500	1 000				<b>7</b>	<b>7,5</b>	0,72	0,77	0,85	0,92	0,98	1,05	1,12	1,19	1,25	1,32	1,37	1,40	1,40	1,37	1,32	1,22	1,12					
1 000	1 500				<b>7</b>	<b>7,5</b>	0,65	0,71	0,77	0,83	0,89	0,95	1,02	1,07	1,13	1,18	1,22	1,25	1,27	1,25	1,20	1,11	1,02					
1 500	1 800				<b>7</b>	<b>7,5</b>	0,57	0,62	0,66	0,71	0,77	0,82	0,88	0,93	0,98	1,02	1,07	1,09	1,10	1,08	1,04	0,97	0,88					
1 800	2 000				<b>9</b>	<b>9,5</b>	0,52	0,56	0,61	0,65	0,70	0,75	0,80	0,84	0,89	0,93	0,98	0,99	1,00	0,99	0,95	0,88	0,80					
2 000	2 500				<b>9</b>	<b>9,5</b>	0,49	0,54	0,58	0,62	0,66	0,71	0,75	0,80	0,84	0,88	0,92	0,94	0,95	0,93	0,89	0,83	0,75					

Feeding rates are indicatives and based on the feed nutritional values. It must be adapted to local conditions and farming goals