




*LAP free : Land Animal Protein free

B-NEO MARIN

Extruded pellet for ongrowing Seabream and Seabass



- Usable for Seabream and Seabass in Organic farming conditions
- Feed with a high DP/DE ratio
- Elevated digestible energy level
- Very high marine products content, selected for their substantial digestibility and palatability level
- Optimized growth performances and FCR (Feed Conversion Ratio)

	B-NEO MARIN
Commercial Ref.	3 / 4 / 5 / 7
Diameter (mm)	3,2 / 4 / 5,5 / 7,5
Presentation	Pellet
Live weight (g)	20 to > 400 g

This fish feed can be used in **Organic farming**, in accordance with the regulation (CE) n°834/2007, (CE) n°889/2008 et (CE) 710/2009. This approach is controlled by FR-BIO 10.

PACKING

20 or 25 kg bag, big bag or bulk
Store feed in a cool and dry place

FLOATING TYPE

Sinking and half-floating

INDICATIVE NUTRITIONAL PROFILE

		B-NEO MARIN
Protein	(%)	43
Fats	(%)	20
Digestible Energy	(MJ/Kg)	18,8
DP / DE	(g/MJ)	20,6
Gross Energy	(MJ/Kg)	21,9
Fibre	(%)	1,8
Ash	(%)	12
Phosphorus	(%)	1,6

VITAMINS INCORPORATION

Vit. A (UI/Kg)	10 000
Vit. D3 (UI/Kg)	1 750
Vit. E (mg/Kg)	200
Vit. C (mg/Kg)	250

SUSTAINABILITY

Here at Le Gouessant, we believe that aquaculture is sustainable if performance, quality and preservation of marine resources are combined, that is why we are Global GAP certified.



FEEDING TABLE

Live Weight (g)		Diameter	Feeding rate (% of Biomass / day) following water temperature																							
From	To		N°	(mm)	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28			
20	50	3	3,2	0,38	0,46	0,55	0,66	0,77	0,92	1,07	1,26	1,45	1,69	1,93	2,20	2,47	2,72	2,98	3,09	3,09	3,09	2,90	2,60			
50	100	4	4	0,29	0,35	0,41	0,50	0,58	0,69	0,80	0,95	1,10	1,27	1,45	1,66	1,86	2,06	2,25	2,33	2,33	2,33	2,19	1,96			
100	150	4	4	0,24	0,29	0,34	0,41	0,48	0,57	0,66	0,78	0,90	1,05	1,20	1,36	1,53	1,69	1,85	1,91	1,91	1,91	1,80	1,62			
150	200	5	5,5	0,21	0,25	0,29	0,35	0,41	0,49	0,58	0,68	0,79	0,91	1,04	1,19	1,33	1,47	1,61	1,67	1,67	1,67	1,57	1,41			
200	300	5	5,5	0,19	0,23	0,27	0,32	0,37	0,45	0,53	0,62	0,72	0,83	0,95	1,08	1,21	1,34	1,47	1,53	1,53	1,53	1,43	1,28			
300	400	5	5,5	0,17	0,20	0,24	0,28	0,33	0,39	0,46	0,54	0,63	0,73	0,83	0,95	1,06	1,17	1,28	1,34	1,34	1,34	1,26	1,13			
400	500	7	7,5	0,14	0,17	0,20	0,24	0,28	0,34	0,40	0,47	0,54	0,63	0,71	0,81	0,91	1,01	1,10	1,14	1,14	1,14	1,07	0,96			
500	600	7	7,5	0,12	0,15	0,18	0,21	0,24	0,29	0,34	0,40	0,47	0,55	0,62	0,71	0,80	0,88	0,96	1,00	1,00	1,00	0,94	0,85			
600	800	7	7,5	0,11	0,13	0,15	0,18	0,22	0,26	0,30	0,36	0,41	0,48	0,55	0,62	0,70	0,77	0,85	0,88	0,88	0,88	0,82	0,73			
800	1 000	7	7,5	0,09	0,11	0,13	0,16	0,19	0,22	0,25	0,30	0,35	0,41	0,47	0,54	0,60	0,66	0,72	0,75	0,75	0,75	0,71	0,63			

Feeding rates are indicatives and based on the feed nutritional values. It must be adapted to local conditions and farming goals