


B-REPRO MARIN



Extruded pellet for Marin Fish Brood stock

- Feed to be used for Marin Fish Brood stock, in Organic conditions
- High marine products content, selected for their high digestibility level
- Enriched with vitamins and omega 3 (EPA & DHA)
- Gametogenesis preparation for eggs yield and survival optimization

| | B-REPRO MARIN |
|------------------|----------------|
| Ref. Commerciale | 7 / 9 |
| Diamètre (mm) | 7,5 / 9,5 |
| Présentation | Pellet |
| Live weight (g) | 400 to 1 000 g |

This fish feed can be used in **Organic farming**, in accordance with the regulation (CE) n°834/2007, (CE) n°889/2008 and (CE) 710/2009. This approach is controlled by FR-BIO 10.

PACKING

20 or 25 kg bag, big bag or bulk
Store feed in a cool and dry place

FLOATING TYPE

Sinking

INDICATIVE NUTRITIONAL PROFILE

| | B-REPRO MARIN |
|----------------------------------|---------------|
| Protein (%) | 51 |
| Fats (%) | 16 |
| Digestible Energy (MJ/Kg) | 18,7 |
| DP / DE (g/MJ) | 25,7 |
| Gross Energy (MJ/Kg) | 21,1 |
| Fibre (%) | 1 |
| Ash (%) | 12 |
| Phosphorus (%) | 1,5 |

VITAMINS INCORPORATION

| | |
|-----------------|--------|
| Vit. A (UI/Kg) | 40 000 |
| Vit. D3 (UI/Kg) | 2 800 |
| Vit. E (mg/Kg) | 600 |
| Vit. C (mg/Kg) | 1 000 |

SUSTAINABILITY

Here at Le Gouessant, we believe that aquaculture is sustainable if performance, quality and preservation of marine resources are combined, that is why we are Global GAP certified.

FEEDING TABLE

| Live Weight (g) | | Feed | P | F | Diameter N° (mm) | Feeding rate (% of Biomass / day) following water temperature | | | | | | | | | | | | | | | | | | |
|-----------------|-------|--------------------------|-----------|-----------|---------------------|---|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| From | To | | | | | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 |
| 400 | 500 | | | | 7 | 0,25 | 0,29 | 0,32 | 0,36 | 0,40 | 0,46 | 0,51 | 0,56 | 0,61 | 0,67 | 0,73 | 0,79 | 0,85 | 0,89 | 0,93 | 0,94 | 0,94 | 0,87 | 0,80 |
| 500 | 600 | | | | 7 | 0,22 | 0,24 | 0,27 | 0,31 | 0,34 | 0,38 | 0,42 | 0,47 | 0,52 | 0,56 | 0,61 | 0,66 | 0,71 | 0,75 | 0,78 | 0,79 | 0,79 | 0,73 | 0,67 |
| 600 | 800 | B-REPRO MARIN | 51 | 16 | 9 | 0,18 | 0,20 | 0,23 | 0,25 | 0,28 | 0,32 | 0,35 | 0,39 | 0,42 | 0,47 | 0,51 | 0,55 | 0,58 | 0,61 | 0,64 | 0,64 | 0,65 | 0,60 | 0,55 |
| 800 | 1 000 | | | | 9 | 0,14 | 0,16 | 0,18 | 0,20 | 0,22 | 0,24 | 0,27 | 0,30 | 0,33 | 0,36 | 0,40 | 0,43 | 0,46 | 0,48 | 0,50 | 0,50 | 0,51 | 0,47 | 0,43 |

Feeding rates are indicatives and based on the feed nutritional values. It must be adapted to local conditions and farming goals