



*LAP free : Land Animal Protein free

B-SPARUS WINTER

Extruded pellet for winter ongrowing
Seabream



- Usable for **Seabream in Organic farming conditions**
- **Feed designed to optimize the resistance of sea bream facing winter conditions.**
- **Very high DP/DE ratio**
- Feed with an **elevated digestible energy level**
- **Highly digestible lipids, 100% marine origin**, formulated without soy
- Supplemented with **B-Glucan, Vitamins and Minerals** to **strengthen seabreams immune system**

	B-SPARUS WINTER
Commerciale Ref.	4 / 5 / 7
Diameter (mm)	4 / 5,5 / 7,5
Presentation	Pellet
Live weight (g)	50 to 1 000 g

This fish feed can be used in **Organic farming**, in accordance with the regulation (CE) n°834/2007, (CE) n°889/2008 et (CE) 710/2009. This approach is controlled by FR-BIO 10.

PACKING

20 and 25 kg bag, big bag or bulk
Store feed in a cool and dry place

FLOATING TYPE

Sinking and half-floating

INDICATIVE NUTRITIONAL PROFILE

		B-SPARUS WINTER
Protein	(%)	47
Fats	(%)	13
Digestible Energy	(MJ/Kg)	17,3
DP / DE	(g/MJ)	25,1
Gross Energy	(MJ/Kg)	19,9
Fibre	(%)	1,3
Ash	(%)	12,4
Phosphorus	(%)	1,7

VITAMINS INCORPORATION

Vit. A (UI/Kg)	12 000
Vit. D3 (UI/Kg)	2 100
Vit. E (mg/Kg)	400
Vit. C (mg/Kg)	500

SUSTAINABILITY

Here at Le Gouessant, we believe that aquaculture is sustainable if performance, quality and preservation of marine resources are combined, that is why we are Global GAP certified.



FEEDING TABLE

Live Weight (g)		Feed	P	F	Diameter (mm)	Feeding rate (% of Biomass / day) following water temperature																						
From	To					10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28				
50	100	B SPARUS WINTER	47	13	4	0,56	0,63	0,71	0,80	0,89	1,00	1,11	1,23	1,35	1,48	1,61	1,74	1,86	1,95	2,05	2,05	2,06	1,91	1,75				
100	150				4	0,47	0,53	0,60	0,67	0,75	0,84	0,92	1,03	1,13	1,24	1,34	1,45	1,56	1,63	1,71	1,71	1,71	1,72	1,59	1,46			
150	200				5	0,41	0,47	0,52	0,59	0,65	0,73	0,82	0,90	0,99	1,09	1,18	1,27	1,36	1,43	1,50	1,50	1,50	1,51	1,39	1,28			
200	300				5	0,36	0,40	0,45	0,51	0,57	0,64	0,71	0,79	0,86	0,95	1,03	1,11	1,19	1,25	1,31	1,31	1,31	1,32	1,23	1,14			
300	400				5	0,31	0,35	0,39	0,44	0,48	0,54	0,60	0,67	0,74	0,82	0,90	0,97	1,03	1,09	1,14	1,14	1,14	1,15	1,07	0,99			
400	500				7	0,27	0,30	0,34	0,38	0,43	0,48	0,53	0,59	0,64	0,71	0,77	0,83	0,89	0,93	0,98	0,98	0,98	0,99	0,91	0,84			
500	600				7	0,23	0,26	0,29	0,32	0,36	0,40	0,44	0,49	0,54	0,59	0,64	0,70	0,75	0,79	0,82	0,83	0,83	0,83	0,77	0,70			
600	800				7	0,19	0,21	0,24	0,27	0,30	0,33	0,37	0,41	0,44	0,49	0,53	0,57	0,61	0,64	0,67	0,68	0,68	0,68	0,63	0,57			
800	1 000				7	0,15	0,17	0,19	0,21	0,23	0,26	0,29	0,32	0,35	0,38	0,42	0,45	0,48	0,50	0,52	0,53	0,53	0,53	0,49	0,45			

Feeding rates are indicatives and based on the feed nutritional values. It must be adapted to local conditions and farming goals