


B-STURGEON RANGE



Extruded pellet for Sturgeons farming

- Feed usable for **Sturgeons in organic farming conditions**
- Range with a high **protein content** and **low lipid level** to **restrict fat accumulation in gonads**
- **Supplemented in vitamin (C and E), minerals and hepatoprotector** to reinforce **gut health**
- Feed **enriched in DHA and omega 3** for **caviar production**

*LAP free : Land Animal Protein free

	B-STURGEON GROWER	B-STURGEON EXTRA	B-STURGEON CAVIAR
Commerciale Ref.	1 / 2 / 3 / 4 / 5 / 7 / 9 / 11 / 13	4 / 5 / 7 / 9	7 / 9 / 11 / 13
Diameter (mm)	1,7 / 2,5 / 3,2 / 4 / 5,5 / 7,5 / 9,5 / 11,5 / 13,5	4 / 5,5 / 7,5 / 9,5	7,5 / 9,5 / 11,5 / 13,5
Presentation	Pellet	Pellet	Pellet
Live weight (g)	5 à 200 g	200 à 3 000 g	3 000 à > 5 000 g

This fish feed for organic shrimp follows the (CE) n°834/2007, (CE) n°889/2008 and (CE) 710/2009 regulations about Organic production. Supervised by FR-BIO 10.

PACKING

25 kg bag, big-bag or bulk
Store feed in a cool and dry place

FLOATING TYPE

Sinking feed (x)

INDICATIVE NUTRITIONAL PROFILE

	B-STURGEON GROWER	B-STURGEON EXTRA	B-STURGEON CAVIAR
Protein (%)	48	45	50
Fats (%)	11	17	11
Gross Energy (MJ/Kg)	19,5	21,1	19,5
Digestible Energy (MJ/Kg)	16,7	18,1	17,1
Starch (%)	14	11,4	13,8
Fibre (%)	1,5	1,9	1
Ash (%)	13,5	10,6	13,8
Phosphorus (%)	1,5	1,4	1,6

VITAMINS INCORPORATION

Vit. A (UI/Kg)	12 000	16 000
Vit. D3 (UI/Kg)	2 100	2 800
Vit. E (mg/Kg)	340	500
Vit. C (mg/Kg)	525	1 000

SUSTAINABILITY

Here at Le Gouessant, we believe that aquaculture is sustainable if performance, quality and preservation of marine resources are combined, that is why we are Global GAP certified.



FEEDING TABLE

Live Weight (g)		Feed	P	F	Diameter		Feeding rate (% of Biomass / day) following water temperature																
From	To				Cial Ref	mm	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29
5	10	B- Sturgeon	48	11	1	1.7	0,81	0,98	1,15	1,27	1,38	1,48	1,58	1,69	1,78	1,85	1,90	1,85	1,75	1,65	1,56	1,40	1,27
10	50				2	2.2	0,72	0,85	1,00	1,11	1,21	1,30	1,40	1,52	1,61	1,68	1,70	1,65	1,57	1,48	1,39	1,28	1,12
50	75				2	2.2	0,63	0,77	0,9	1,00	1,10	1,18	1,26	1,37	1,47	1,50	1,52	1,46	1,4	1,33	1,26	1,15	1,04
75	200				3	3.2	0,53	0,69	0,84	0,94	1,04	1,10	1,15	1,21	1,26	1,28	1,30	1,25	1,20	1,12	1,04	0,94	0,84
200	500	B-Sturgeon Extra	45	17	4	4	0,48	0,62	0,75	0,83	0,90	0,95	1,00	1,05	1,10	1,12	1,14	1,07	1,00	0,95	0,90	0,80	0,70
500	1000				4	4	0,40	0,52	0,63	0,72	0,80	0,85	0,90	0,93	0,96	0,98	1,00	0,93	0,86	0,79	0,72	0,61	0,50
1000	2000				5	5.5	0,32	0,36	0,40	0,50	0,60	0,65	0,70	0,73	0,75	0,78	0,80	0,73	0,65	0,58	0,50	0,43	0,35
2000	3000				7	7.5	0,13	0,17	0,20	0,25	0,30	0,33	0,36	0,40	0,43	0,44	0,45	0,41	0,36	0,33	0,30	0,25	0,20
3000	5000	B-Sturgeon Caviar	50	11	7	7.5	0,08	0,10	0,12	0,15	0,17	0,19	0,20	0,22	0,23	0,24	0,25	0,24	0,22	0,20	0,17	0,14	0,10
> 5000					9 / 11	9.5 / 11.5	0,06	0,08	0,10	0,12	0,13	0,15	0,16	0,17	0,18	0,19	0,20	0,19	0,18	0,16	0,13	0,11	0,08

Feeding rates are indicatives and based on the feed nutritional values. It must be adapted to local conditions and farming goals