

# ORGANIC HATCHERY RANGE : B-SUPRA – B-START MARIN



## Extruded feed for **Marin fish fingerlings**

- Feed usable for juveniles in **Organic** farming conditions
- **Progressive protein and lipid profile**, according to the nutritional need of each life stage
- **High DP/DE ratio**
- High **marine products proportion**
- **Supplemented feed in Vitamins and Minerals**, to strengthen the **immune system**

Commercial Ref.	B-SUPRA			B-START MARIN	
	M1 / M2	AL2 / AL3	AL4	1	2 / 3
Diameter (mm)	0,4-0,7 / 0,7-1	0,8 / 1,1	1,4	1,7	2,5 / 3,3
Presentation	Crumble	Pellet	Pellet	Pellet	Pellet
Live weight (g)	0,1 to 1,5 g	1,5 to 2,5 g	2,5 to 5 g	5 to 10 g	10 to 50 g

This fish feed can be used in **Organic farming**, in accordance with the regulation (CE) n°834/2007, (CE) n°889/2008 et (CE) 710/2009. This approach is controlled by FR-BIO 10.

### PACKING

**B-SUPRA** : 20 kg bag or 10 kg bucket  
**B-START MARIN**: 25 kg (20 kg for half-floating)

Store feed in a cool and dry place

### FLOATING TYPE

**B-SUPRA**: Sinking (slowly)  
**B-START MARIN**: Sinking or half-floating

### INDICATIVE PROFILE NUTRITIONAL

		B-SUPRA		B-START MARIN	
		M1 / M2 / AL2 / AL3	AL4	1	2 / 3
<b>Protein</b>	(%)	<b>55</b>	<b>52</b>	<b>47</b>	<b>45</b>
<b>Fats</b>	(%)	<b>13</b>	<b>17</b>	<b>15</b>	<b>17</b>
<b>Digestible Energy (DE)</b>	(MJ/Kg)	<b>18,3</b>	<b>19,2</b>	<b>17,8</b>	<b>18,2</b>
<b>DP / DE</b>	(g/MJ)	29,1	25,1	24	22,7
<b>Gross Energy</b>	(MJ/Kg)	20,5	21,8	20,7	21
<b>Fibres</b>	(%)	0,5	1,1	1,5	1,3
<b>Ash</b>	(%)	14	11,9	12,3	13
<b>Phosphorus</b>	(%)	1,8	1,5	1,7	1,7

### VITAMINS INCORPORATION

Vit. A (UI/Kg)	12 000	10 000
Vit. D3 (UI/Kg)	2 100	1 750
Vit. E (mg/Kg)	400	200
Vit. C (mg/Kg)	1 000	250

### SUSTAINABILITY

Here at Le Gouessant, we believe that aquaculture is sustainable if performance, quality and preservation of marine resources are combined, that is why we are Global GAP certified.



# FEEDING TABLE

Live Weight (g)		Feed	P	F	Diameter		Feeding rate (% of biomass / day) following temperature (°C)																
From	To				N°	(mm)	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28		
0,1	0,5	<b>B SUPRA</b>	55	13	M1	0,4-0,7	2,35	2,62	2,90	3,20	3,51	3,84	4,19	4,55	4,93	5,33	5,57	5,60	5,33	4,90	4,47		
0,5	1,5					0,7-1	2,09	2,33	2,58	2,85	3,13	3,43	3,74	4,06	4,40	4,76	5,04	5,08	4,76	4,22	3,89	3,55	
1,5	2,5					1,2	1,86	2,07	2,29	2,53	2,78	3,04	3,32	3,61	3,91	4,23	4,51	4,52	4,23	3,75	3,42	3,08	2,76
2,5	5					1,4	1,56	1,73	1,92	2,12	2,33	2,55	2,79	3,03	3,29	3,55	3,72	3,75	3,42	3,06	2,71	2,37	2,03
5	10	<b>B START MARIN</b>	47	15	1	1,7	1,63	1,82	2,01	2,22	2,44	2,67	2,91	3,16	3,42	3,68	3,87	3,90	3,71	3,42	3,12		
10	15					2,5	1,44	1,61	1,78	1,97	2,16	2,37	2,58	2,80	3,03	3,25	3,42	3,44	3,29	3,02	2,76		
15	20					2	1,34	1,50	1,66	1,83	2,01	2,20	2,40	2,60	2,82	3,02	3,17	3,20	3,06	2,82	2,57		
20	25					3	1,27	1,41	1,56	1,73	1,90	2,08	2,27	2,46	2,66	2,85	2,99	3,02	2,89	2,66	2,43		
25	30	<b>B START MARIN</b>	45	17	3	3,2	1,21	1,34	1,49	1,65	1,81	1,98	2,16	2,34	2,53	2,71	2,85	2,87	2,75	2,54	2,32		
30	35					3	1,16	1,29	1,43	1,58	1,73	1,90	2,07	2,25	2,43	2,60	2,73	2,75	2,64	2,43	2,22		
35	40					3	1,11	1,24	1,38	1,52	1,67	1,83	2,00	2,16	2,34	2,50	2,62	2,64	2,55	2,34	2,14		
40	45					3	1,08	1,20	1,33	1,47	1,62	1,77	1,93	2,09	2,26	2,42	2,53	2,55	2,46	2,27	2,07		
45	50	<b>B START MARIN</b>	45	17	3	3,2	1,04	1,16	1,29	1,43	1,57	1,72	1,87	2,03	2,19	2,34	2,45	2,47	2,39	2,20	2,01		

Feeding rates are indicatives and based on the feed nutritional values. It must be adapted to local conditions and farming goals