




CATFISH RANGE

Extruded feed for Catfish



- Floating pellets with a high water stability
- Supplemented fingerling feed to strengthen their immune system
- Use of essential oils with anti-bacterial properties, to preventive purpose, to limite pathological pressure
- Inclusion of marine products in order to increase palatability and optimize feed efficiency
- High digestibility feed range

*LAP free : Land Animal Protein free

	NEO SUPRA		CATFISH STARTER	CATFISH GROWER	CATFISH FINISHER
Commerciale Ref.	AL0 / AL1 / AL2	AL2 / AL3	1 / 2	3 / 4 / 5	5 / 7 / 9
Diameter (mm)	0,15-0,4 / 0,4-0,7 / 0,7-1	0,8 / 1,2	1,7 / 2,5	3,2 / 4 / 5,5	5,5 / 7,5 / 9,5
Presentation	Crumble	Pellet	Pellet	Pellet	Pellet
Live weight (g)	< 0,5 to 5 g		5 to 50 g	50 to 1 000 g	1 000 to >1 500 g

PACKING

NEO SUPRA : 20 kg bag (or 10 kg bucket)
CATFISH STARTER, GROWER and FINISHER : 15 kg bag
 Store feed in a cool and dry place .

FLOATING TYPE

NEO SUPRA: Half-floating (Y)
CATFISH STARTER, GROWER and FINISHER : Floating (Z)

INDICATIVE PROFILE NUTRITIONAL

	NEO SUPRA	CATFISH STARTER	CATFISH GROWER	CATFISH FINISHER
Protein (%)	58	46	42	38
Fats (%)	12	10	11	12
Gross Energy (MJ/Kg)	21,2	19,5	19,7	19,7
Starch (%)	11,3	15,6	14,5	18,8
Fibre (%)	0,5	3	3,8	3,7
Ash (%)	9	9,2	7,3	6,5
Phosphorus (%)	1,5	1,3	1	0,9

VITAMINS INCORPORATION

Vit. A (UI/Kg)	12 000	8 000
Vit. D3 (UI/Kg)	2 100	1 400
Vit. E (mg/Kg)	440	160
Vit. C (mg/Kg)	1 000	150

SUSTAINABILITY

Here at Le Gouessant, we believe that aquaculture is sustainable if performance, quality and preservation of marine resources are combined, that is why we are Global GAP certified.



FEEDING TABLE

Live Weight (g)		Feed	P	F	Diameter	Feeding rate (% of Biomass / day) following water temperature																			
From	To					N°	(mm)	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33
< 0,5		NEO SUPRA	58	13	AL0	Ad libitum																			
					AL1	Ad libitum																			
					AL2	Ad libitum																			
2,5		5	46	10	AL3	1,2	1,21	1,82	2,42	3,23	4,04	4,91	5,77	6,49	7,21	7,61	8,02	8,23	8,44	8,31	8,18	7,98	7,78	7,39	7,00
5	10				1	1,7	1,09	1,64	2,18	2,91	3,64	4,42	5,19	5,84	6,49	6,85	7,22	7,41	7,59	7,48	7,37	7,18	7,00	6,65	6,30
10		50	42	11	2	2,5	0,82	1,23	1,64	2,18	2,73	3,31	3,90	4,38	4,87	5,14	5,41	5,55	5,70	5,61	5,53	5,39	5,25	4,99	4,72
50	100				3	3,2	0,64	0,95	1,27	1,70	2,12	2,58	3,03	3,41	3,79	4,00	4,21	4,32	4,43	4,36	4,30	4,19	4,08	3,88	3,67
100		150	42	11	5	5,5	0,41	0,61	0,82	1,09	1,36	1,66	1,95	2,19	2,44	2,57	2,71	2,78	2,85	2,81	2,76	2,69	2,62	2,49	2,36
100	150				5	5,5	0,24	0,35	0,47	0,63	0,79	0,96	1,13	1,27	1,41	1,49	1,56	1,60	1,65	1,62	1,60	1,56	1,52	1,44	1,36
300		500	42	11	5	5,5	0,35	0,53	0,71	0,94	1,17	1,43	1,68	1,88	2,09	2,22	2,33	2,39	2,46	2,42	2,38	2,33	2,26	2,15	2,04
300	500				5	5,5	0,38	0,57	0,77	1,02	1,27	1,54	1,82	2,04	2,26	2,39	2,51	2,59	2,66	2,61	2,57	2,51	2,44	2,32	2,20
500		800	38	12	7	7,5	0,29	0,44	0,59	0,79	0,98	1,19	1,40	1,58	1,75	1,85	1,94	2,00	2,05	2,02	1,99	1,94	1,89	1,79	1,70
500	800				7	7,5	0,26	0,38	0,51	0,68	0,85	1,04	1,22	1,37	1,52	1,61	1,69	1,73	1,78	1,75	1,73	1,69	1,64	1,56	1,47
800		1000	38	12	7	7,5	0,20	0,30	0,40	0,54	0,67	0,81	0,96	1,08	1,20	1,26	1,33	1,37	1,40	1,38	1,36	1,33	1,29	1,23	1,16
800	1000				7	7,5	0,16	0,25	0,33	0,44	0,55	0,67	0,78	0,88	0,98	1,03	1,09	1,12	1,15	1,13	1,11	1,08	1,06	1,00	0,95
> 1500		1500	38	12	7	7,5	0,16	0,25	0,33	0,44	0,55	0,67	0,78	0,88	0,98	1,03	1,09	1,12	1,15	1,13	1,11	1,08	1,00	0,95	
> 1500	1500				7	7,5	0,16	0,25	0,33	0,44	0,55	0,67	0,78	0,88	0,98	1,03	1,09	1,12	1,15	1,13	1,11	1,08	1,06	1,00	0,95

Feeding rates are indicatives and based on the feed nutritional values. It must be adapted to local conditions and farming goals