



*LAP free : Land Animal Protein free

NEO SPARUS WINTER



Extruded pellet for ongrowing **SEABREAM** in winter condition

- Feed designed to optimize the resistance of sea bream facing winter conditions.
- Very high DP/DE ratio, feed with an elevated digestible energy level
- Highly digestible lipids, 100% marine origin, formulated without soya
- Supplemented with β -Glucan, Hepathoprotector, Vitamins and Minerals to strengthen Seabream immune system during critical times
- Use of essential oils with antibacterial properties, for preventive purpose, to limit pathological pressure

| | NEO SPARUS WINTER |
|------------------|-------------------|
| Commerciale Ref. | 4 / 5 / 7 |
| Diameter (mm) | 4 / 5,5 / 7,5 |
| Presentation | Pellet |
| Live weight (g) | 50 to > 400 g |

PACKING

20 or 25 kg bag, big bag or bulk

Store feed in a cool and dry place

FLOATING TYPE

Sinking and half-floating

INDICATIVE PROFILE NUTRITIONAL

| | | NEO SPARUS WINTER |
|-------------------|---------|-------------------|
| Protein | (%) | 47 |
| Fats | (%) | 13 |
| Digestible Energy | (MJ/Kg) | 17,6 |
| DP / DE | (g/MJ) | 24,7 |
| Gross Energy | (MJ/Kg) | 20,4 |
| Fibre | (%) | 2,5 |
| Ash | (%) | 8,5 |
| Phosphorus | (%) | 1,2 |

VITAMINS INCORPORATION

| | |
|-----------------|--------|
| Vit. A (UI/Kg) | 12 000 |
| Vit. D3 (UI/Kg) | 2 100 |
| Vit. E (mg/Kg) | 400 |
| Vit. C (mg/Kg) | 500 |

SUSTAINABILITY

Here at Le Gouessant, we believe that aquaculture is sustainable if performance, quality and preservation of marine resources are combined, that is why we are Global GAP certified.



FEEDING TABLE

| Live Weight (g) | | Feed | P | F | Diameter | Feeding rate (% of Biomass / day) following water temperature | | | | | | | | | | | | | | | | | | | | | | |
|-----------------|-------|------|---|---|----------|---------------------------------------------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|----|--|--|
| From | To | | | | | N° | (mm) | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | | |
| 50 | 100 | | | | 4 | 4 | 0,56 | 0,63 | 0,71 | 0,80 | 0,89 | 1,00 | 1,11 | 1,23 | 1,35 | 1,48 | 1,61 | 1,74 | 1,86 | 1,95 | 2,05 | 2,05 | 2,06 | 1,91 | 1,75 | | | |
| 100 | 150 | | | | 4 | 4 | 0,47 | 0,53 | 0,60 | 0,67 | 0,75 | 0,84 | 0,92 | 1,03 | 1,13 | 1,24 | 1,34 | 1,45 | 1,56 | 1,63 | 1,71 | 1,71 | 1,72 | 1,59 | 1,46 | | | |
| 150 | 200 | | | | 5 | 5,5 | 0,41 | 0,47 | 0,52 | 0,59 | 0,65 | 0,73 | 0,82 | 0,90 | 0,99 | 1,09 | 1,18 | 1,27 | 1,36 | 1,43 | 1,50 | 1,50 | 1,51 | 1,39 | 1,28 | | | |
| 200 | 300 | | | | 5 | 5,5 | 0,36 | 0,40 | 0,45 | 0,51 | 0,57 | 0,64 | 0,71 | 0,79 | 0,86 | 0,95 | 1,03 | 1,11 | 1,19 | 1,25 | 1,31 | 1,31 | 1,32 | 1,23 | 1,14 | | | |
| 300 | 400 | | | | 5 | 5,5 | 0,31 | 0,35 | 0,39 | 0,44 | 0,48 | 0,54 | 0,60 | 0,67 | 0,74 | 0,82 | 0,90 | 0,97 | 1,03 | 1,09 | 1,14 | 1,14 | 1,15 | 1,07 | 0,99 | | | |
| 400 | 500 | | | | 7 | 7,5 | 0,27 | 0,30 | 0,34 | 0,38 | 0,43 | 0,48 | 0,53 | 0,59 | 0,64 | 0,71 | 0,77 | 0,83 | 0,89 | 0,93 | 0,98 | 0,98 | 0,99 | 0,91 | 0,84 | | | |
| 500 | 600 | | | | 7 | 7,5 | 0,23 | 0,26 | 0,29 | 0,32 | 0,36 | 0,40 | 0,44 | 0,49 | 0,54 | 0,59 | 0,64 | 0,70 | 0,75 | 0,79 | 0,82 | 0,83 | 0,83 | 0,77 | 0,70 | | | |
| 600 | 800 | | | | 7 | 7,5 | 0,19 | 0,21 | 0,24 | 0,27 | 0,30 | 0,33 | 0,37 | 0,41 | 0,44 | 0,49 | 0,53 | 0,57 | 0,61 | 0,64 | 0,67 | 0,68 | 0,68 | 0,63 | 0,57 | | | |
| 800 | 1 000 | | | | 7 | 7,5 | 0,15 | 0,17 | 0,19 | 0,21 | 0,23 | 0,26 | 0,29 | 0,32 | 0,35 | 0,38 | 0,42 | 0,45 | 0,48 | 0,50 | 0,52 | 0,53 | 0,53 | 0,49 | 0,45 | | | |

Feeding rates are indicatives and based on the feed nutritional values. It must be adapted to local conditions and farming goals