



## NEO SPARUS

### Extruded pellet for ongrowing Seabream



- Feed with a very **high DP/DE ratio**, and an **elevated digestible energy** level
- Very **high marine products** content, selected for their **substantial digestibility and palatability level**
- Feed supplemented with **EPA & DHA**
- **Supplemented** with several pigments (**Spirulina, Astaxanthin, Xanthophylls and Carotenoids**) to intensify seabreams **pigmentation**

\*LAP free : Land Animal Protein free

Commerciale Ref.	NEO SPARUS	
	3 / 4	5 / 7
Diameter (mm)	3,2 / 4	5,5 / 7,5
Presentation	Pellet	Pellet
Live weight (g)	20 to 150 g	150 to > 800 g

#### PACKING

15, 20 or 25 kg bag, big bag or bulk

Store feed in a cool and dry place.

#### FLOATING TYPE

Sinking, half-floating or floating

#### INDICATIVE NUTRITIONAL PROFILE

		NEO SPARUS	
<b>Protein</b>	<b>(%)</b>	<b>45</b>	<b>43</b>
<b>Fats</b>	<b>(%)</b>	<b>18</b>	<b>19</b>
<b>Digestible Energy</b>	<b>(MJ/Kg)</b>	<b>18,5</b>	<b>18,5</b>
<b>DP / DE</b>	<b>(g/MJ)</b>	21,4	21
<b>Gross Energy</b>	<b>(%)</b>	2	3
<b>Fibre</b>	<b>(%)</b>	8	8
<b>Ash</b>	<b>(%)</b>	14	13
<b>Phosphorus</b>	<b>(%)</b>	0,9	0,9

#### VITAMINS INCORPORATION

Vit. A (UI/Kg)	10 000	12 000
Vit. D3 (UI/Kg)	1 750	2 100
Vit. E (mg/Kg)	200	200
Vit. C (mg/Kg)	150	150

#### SUSTAINABILITY

Here at Le Gouessant, we believe that aquaculture is sustainable if performance, quality and preservation of marine resources are combined, that is why we are Global GAP certified.



## FEEDING TABLE

Live Weight (g)		Feed	P	F	Diameter	Feeding rate (% of Biomass / day) following water temperature																					
From	To					N°	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28		
20	50		45	18	3	0,70	0,79	0,89	1,00	1,12	1,25	1,39	1,54	1,69	1,85	2,01	2,17	2,32	2,43	2,54	2,56	2,57	2,38	2,18			
50	100					4	0,53	0,60	0,68	0,76	0,85	0,95	1,06	1,17	1,29	1,41	1,53	1,65	1,77	1,86	1,95	1,95	1,95	1,96	1,81	1,66	
100	150				4	0,45	0,51	0,57	0,64	0,71	0,79	0,88	0,98	1,08	1,18	1,28	1,38	1,48	1,55	1,63	1,63	1,64	1,51	1,39			
150	200	NEO SPARUS	43	19	5	0,39	0,44	0,49	0,56	0,62	0,70	0,78	0,86	0,94	1,03	1,12	1,21	1,30	1,36	1,43	1,43	1,43	1,32	1,21			
200	300					5	0,34	0,38	0,43	0,48	0,54	0,61	0,68	0,75	0,82	0,90	0,98	1,06	1,14	1,19	1,24	1,25	1,25	1,25	1,17	1,08	
300	400					5	0,30	0,34	0,37	0,42	0,46	0,52	0,57	0,64	0,70	0,78	0,86	0,92	0,98	1,03	1,08	1,08	1,09	1,09	1,09	1,02	0,94
400	500					7	0,25	0,29	0,32	0,36	0,40	0,46	0,51	0,56	0,61	0,67	0,73	0,79	0,85	0,89	0,93	0,93	0,94	0,94	0,94	0,87	0,80
500	600				7	0,22	0,24	0,27	0,31	0,34	0,38	0,42	0,47	0,52	0,56	0,61	0,66	0,71	0,75	0,78	0,79	0,79	0,73	0,67			
600	800				7	0,18	0,20	0,23	0,25	0,28	0,32	0,35	0,39	0,42	0,47	0,51	0,55	0,58	0,61	0,64	0,64	0,65	0,60	0,55			
800	1000				7	0,14	0,16	0,18	0,20	0,22	0,24	0,27	0,30	0,33	0,36	0,40	0,43	0,46	0,48	0,50	0,50	0,51	0,47	0,43			

Feeding rates are indicatives and based on the feed nutritional values. It must be adapted to local conditions and farming goals