



RAS HATCHERY RANGE :

Extruded pellet for **Marin Fish Fingerlings** in **Recirculated Aquaculture System**

- **Progressive protein and lipid profile**, according to the nutritional need of each life stage, high DP/DE ratio
- **High marine products proportion**
- **Supplemented feed to strengthen juveniles immune system: MOS, β -Glucan, Vitamins and Minerals**
- **Use of essential oils with antibacterial properties**, for preventive purpose, to limit pathological pressure
- **Adapted range for Recirculated Aquaculture System, supplemented with additive to densify faeces, making them easier to collect.**



*LAP free : Land Animal Protein free

	NEO SUPRA MARIN	NEO SUPRA LOOP		NEO START LOOP	
Commerciale Ref.	M1 / M2	AL2 / AL3	AL4	1	2 / 3
Diameter (mm)	0,4-0,7 / 0,7-1	0,8 / 1,1	1,4	1,7	2,5 / 3,2
Presentation	Crumble	Pellet	Pellet	Pellet	Pellet
Live weight (g)	0,1 to 1,5 g	1,5 to 2,5 g	2,5 to 5 g	5 to 10 g	10 to 50 g

PACKING

NEO SUPRA LOOP: 20 kg bag or 10 kg bucket

NEO START LOOP : 25 kg bag

Store feed in a cool and dry place.

FLOATING TYPE

NEO SUPRA LOOP: Sinking (slowly)

NEO START LOOP: Sinking

INDICATIVE NUTRITIONAL PROFILE

		NEO SUPRA MARIN	NEO SUPRA LOOP		NEO START LOOP	
		M1 / M2	AL2 / AL3	AL4	1	2 / 3
Protein	(%)	58	58	55	52	47
Fats	(%)	13	13	16	17	18
Digestible Energy	(MJ/Kg)	18,8	18,8	19,5	19,5	19,2
DP / DE	(g/MJ)	29,5	29,5	27,1	25,5	23
Gross Energy	(MJ/Kg)	21,1	21,1	21,8	22	21,9
Fibre	(%)	0,3	0,3	0,5	1	1,5
Ash	(%)	10	10	9	9	8
Phosphorus	(%)	1,5	1,5	1,5	1,5	1,2

VITAMINS INCORPORATION

Vit. A (UI/Kg)	12 000	10 000
Vit. D3 (UI/Kg)	2 100	1 750
Vit. E (mg/Kg)	400	200
Vit. C (mg/Kg)	1 000	250

SUSTAINABILITY

Here at Le Gouessant, we believe that aquaculture is sustainable if performance, quality and preservation of marine resources are combined, that is why we are Global GAP certified.



FEEDING TABLE

Live Weight (g)		Feed	P	F	Diameter		Feeding rate (% of biomass / day) following temperature (°C)														
From	To				N°	(mm)	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
0,1	0,5	NEO SUPRA MARIN	58	13	M1	0,4-0,7	2,55	2,84	3,15	3,47	3,81	4,17	4,55	4,94	5,35	5,78	6,09	6,17	5,78	5,31	4,84
0,5	1,5				M2	0,7-1	2,04	2,27	2,51	2,77	3,05	3,34	3,64	3,96	4,29	4,63	4,91	4,95	4,63	4,26	3,89
0,5	1,5	NEO SUPRA LOOP	58	13	AL2	0,7-1	2,04	2,27	2,51	2,77	3,05	3,34	3,64	3,96	4,29	4,63	4,91	4,95	4,63	4,26	3,89
1,5	2,5				AL3	1,2	1,806	2,011	2,229	2,459	2,703	2,959	3,228	3,51	3,805	4,113	4,394	4,404	4,111	3,783	3,455
2,5	5				AL4	1,4	1,533	1,707	1,892	2,088	2,295	2,514	2,743	2,984	3,236	3,499	3,662	3,692	3,495	3,218	2,942
5	10	NEO START LOOP	47	18	1	1,7	1,488	1,658	1,837	2,029	2,229	2,44	2,66	2,887	3,125	3,362	3,531	3,562	3,389	3,118	2,848
10	15				2	2,5	1,37	1,52	1,69	1,86	2,05	2,24	2,44	2,65	2,87	3,08	3,24	3,26	3,12	2,87	2,62
15	20				2	2,5	1,27	1,42	1,57	1,73	1,91	2,09	2,27	2,47	2,67	2,87	3,01	3,03	2,90	2,67	2,44
20	25				3	3,2	1,20	1,34	1,48	1,64	1,80	1,97	2,15	2,33	2,52	2,70	2,84	2,86	2,74	2,52	2,31
25	30				3	3,2	1,14	1,27	1,41	1,56	1,71	1,88	2,05	2,22	2,40	2,57	2,70	2,72	2,61	2,40	2,20
30	35				3	3,2	1,10	1,22	1,35	1,50	1,64	1,80	1,96	2,13	2,30	2,46	2,58	2,60	2,50	2,31	2,11
35	40	3	3,2	1,05	1,18	1,30	1,44	1,58	1,73	1,89	2,05	2,22	2,37	2,49	2,51	2,41	2,22	2,03			
40	45	3	3,2	1,02	1,14	1,26	1,39	1,53	1,68	1,83	1,98	2,14	2,29	2,40	2,42	2,33	2,15	1,97			
45	50	3	3,2	0,99	1,10	1,22	1,35	1,48	1,63	1,77	1,92	2,07	2,22	2,33	2,34	2,26	2,08	1,91			

Feeding rates are indicatives and based on the feed nutritional values. It must be adapted to local conditions and farming goals