




*LAP free : Land Animal Protein free

HATCHERY RANGE : NEO SUPRA - NEO START



Extruded pellet for **Marin fish fingerlings**

- **Progressive protein and lipid profile**, according to the nutritional need of each life stage, high DP/DE ratio
- **High marine products proportion**
- **Supplemented feed to strengthen Trout juveniles immune system: MOS, β -Glucan, Vitamins and Minerals**
- **Use of essential oils with antibacterial properties**, for preventive purpose, to **limit pathological pressure**

Commerciale Ref.	NEO SUPRA MARIN			NEO START	
	M1 / M2	AL2 / AL3	AL4	1	2 / 3
Diameter (mm)	0,4-0,7 / 0,7-1	0,8 / 1,1	1,4	1,7	2,5 / 3,2
Presentation	Crumble	Pellet	Pellet	Pellet	Pellet
Live weight (g)	0,1 à 1,5 g	1,5 à 2,5 g	2,5 à 5 g	5 à 10 g	10 à 50 g

PACKING

NEO SUPRA MARIN : 20 Kg bag or 10 Kg bucket
NEO START: 25 Kg bag (20 kg if half-floating)

Store feed in a cool and dry place

FLOATING TYPE

NEO SUPRA MARIN: sinking (slowly)
NEO START : sinking and half-floating

INDICATIVE NUTRITIONAL PROFILE

		NEO SUPRA MARIN		NEO START	
		M1 / M2 / AL2	AL3 / AL4	1	2 / 3
Protéine	(%)	58		52	47
Lipides	(%)	13		17	18
Energie Digestible (ED)	(MJ/Kg)	19		19,2	19
PD / ED	(g/MJ)	29		25,5	23
Energie Brute	(MJ/Kg)	21,2		21,7	21,4
Cellulose	(%)	0,5		1	1,5
Cendres	(%)	9		9	8
Phosphore	(%)	1,5		1,5	1

VITAMINS INCORPORATION

Vit. A (UI/Kg)	12 000	10 000
Vit. D3 (UI/Kg)	2 100	1 750
Vit. E (mg/Kg)	400	200
Vit. C (mg/Kg)	1 000	250

SUSTAINABILITY

Here at Le Gouessant, we believe that aquaculture is sustainable if performance, quality and preservation of marine resources are combined, that is why we are Global GAP certified.



FEEDING TABLE

Live Weight (g)		Feed	P	F	Diameter	Feeding rate (% of biomass / day) following temperature (°C)															
From	To					N°	(mm)	14	15	16	17	18	19	20	21	22	23	24	25	26	27
0,1	0,5	NEO SUPRA MARIN	58	13	M1	0,4-0,7	2,26	2,52	2,79	3,08	3,38	3,70	4,04	4,39	4,75	5,13	5,37	5,39	5,14	4,72	4,30
0,5	1,5					0,7-1	2,02	2,25	2,49	2,75	3,02	3,30	3,60	3,91	4,24	4,58	4,86	4,89	4,58	4,22	3,85
1,5	2,5					1,2	1,79	1,99	2,21	2,43	2,67	2,93	3,19	3,47	3,76	4,07	4,35	4,36	4,07	3,74	3,42
2,5	5					1,4	1,57	1,75	1,94	2,14	2,36	2,58	2,82	3,06	3,32	3,59	3,76	3,79	3,59	3,30	3,02
5	10	NEO START	52	17	1	1,7	1,51	1,68	1,87	2,06	2,26	2,48	2,70	2,93	3,17	3,41	3,59	3,62	3,44	3,17	2,89
10	15					2,5	1,38	1,54	1,71	1,88	2,07	2,27	2,47	2,68	2,90	3,12	3,27	3,30	3,15	2,90	2,65
15	20					2,5	1,28	1,43	1,59	1,75	1,93	2,11	2,30	2,49	2,70	2,90	3,04	3,06	2,93	2,70	2,47
20	25					3,2	1,21	1,35	1,50	1,66	1,82	1,99	2,17	2,35	2,55	2,73	2,87	2,89	2,77	2,55	2,33
25	30	NEO START	47	18	3	3,2	1,15	1,29	1,43	1,58	1,73	1,90	2,07	2,24	2,43	2,60	2,73	2,75	2,64	2,43	2,22
30	35					3,2	1,11	1,24	1,37	1,51	1,66	1,82	1,98	2,15	2,32	2,49	2,61	2,63	2,53	2,33	2,13
35	40					3,2	1,07	1,19	1,32	1,46	1,60	1,75	1,91	2,07	2,24	2,40	2,51	2,53	2,44	2,25	2,05
40	45					3,2	1,03	1,15	1,27	1,41	1,55	1,69	1,85	2,00	2,16	2,31	2,43	2,44	2,36	2,17	1,99
45	50	NEO START	47	18	3	3,2	1,00	1,11	1,24	1,37	1,50	1,64	1,79	1,94	2,10	2,24	2,35	2,37	2,29	2,11	1,93

Feeding rates are indicatives and based on the feed nutritional values. It must be adapted to local conditions and farming goals