


# HATCHERY RANGE: NEO SUPRA - NEO START



## Extruded pellet for Sturgeon fry

- **Progressive evolution of the protein and lipid profil**, according to the nutritional need of each stage, high DP/DE ratio
- **High proportion of marine products**
- **Supplemented to strengthen the immune system** of alevins: **MOS, B-Glucan, Vitamins and Minerals**
- **Use of essential oils with antibacterial properties**, for preventive purpose, to limit health pressure

\*LAP free : Land Animal Protein free

Commerciale Ref.	NEO SUPRA				NEO START	
	M00 / M0	M1 / M2	AL2 / AL3	AL4	1	2 / 3
Diameter (mm)	<0,15 / 0,15-0,4	0,4-0,7 / 0,7-1	0,8 / 1,1	1,4	1,7	2,5 / 3,2
Presentation	Miette	Miette	Granulé	Granulé	Granulé	Granulé
Live weight (g)	< 0,2	0,2 à 1,5 g	1,5 à 2,5 g	2,5 à 5 g	5 à 10 g	10 à 50 g

### PACKING

**NEO SUPRA** : 20 Kg bag or 10 Kg bucket

**NEO START**: 25 Kg bag

*Store feed in a cool and dry place*

### FLOATING TYPE

**NEO SUPRA** : sinking (slowly)

**NEO START** : sinking

### INDICATIVE PROFILE NUTRITIONAL

		NEO SUPRA		NEO START	
				1	2 / 3
<b>Protein</b>	<b>(%)</b>	<b>58</b>		<b>52</b>	<b>47</b>
<b>Fats</b>	<b>(%)</b>	<b>13</b>		<b>17</b>	<b>18</b>
<b>Digestible Energy</b>	<b>(MJ/Kg)</b>	<b>19</b>		<b>19,2</b>	<b>19</b>
<b>DP / DE</b>	<b>(g/MJ)</b>	29		25,5	23
<b>Gross Energy</b>	<b>(MJ/Kg)</b>	21,2		21,7	21,4
<b>Fibre</b>	<b>(%)</b>	0,5		1	1,5
<b>Ash</b>	<b>(%)</b>	9		9	8
<b>Phosphorus</b>	<b>(%)</b>	1,5		1,5	1

### VITAMINS INCORPORATION

Vit. A (UI/Kg)	12 000	10 000
Vit. D3 (UI/Kg)	2 100	1 750
Vit. E (mg/Kg)	440	200
Vit. C (mg/Kg)	1 000	150

### SUSTAINABILITY

Here at Le Gouessant, we believe that aquaculture is sustainable if performance, quality and preservation of marine resources are combined, that is why we are Global GAP certified.



## FEEDING TABLE

Live Weight (g)		Feed	P	F	Diameter	Feeding rate (% of Biomass / day) following water temperature																	
From	To					N°	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27
0,1	0,5				M1	0,4-0,7	2,26	2,50	2,77	2,98	3,20	3,50	3,80	4,10	4,40	4,74	5,05	5,44	5,53	4,93	3,90	2,74	1,87
0,5	1,5	NEO	58	13	M2	0,7-1	1,95	2,13	2,30	2,50	2,74	2,98	3,27	3,55	3,84	4,08	4,37	4,66	4,80	4,23	3,46	2,66	1,87
1,5	2,5	SUPRA			AL3	1,2	1,74	1,90	2,05	2,20	2,42	2,62	2,87	3,13	3,40	3,58	3,82	4,00	4,10	3,68	3,10	2,50	1,87
2,5	5				AL4	1,4	1,49	1,63	1,77	1,91	2,05	2,23	2,42	2,60	2,79	2,98	3,16	3,29	3,37	3,21	2,77	2,32	1,84
5	10	NEO	52	17	1	1,7	1,26	1,35	1,46	1,56	1,65	1,74	1,84	1,95	2,05	2,15	2,24	2,30	2,38	2,38	2,32	2,10	1,77
10	50	START	47	18	2	2,5	1,16	1,25	1,35	1,45	1,54	1,64	1,74	1,84	1,93	2,02	2,10	2,15	2,20	2,20	2,16	1,98	1,72

Feeding rates are indicatives and based on the feed nutritional values. It must be adapted to local conditions and farming goals