

PERCID RANGE

Extruded pellet for **Perch, Pike Perch and others carnivorous freshwater fish**



- **Supplemented fingerling feed to strengthen their immune system**
- **Use of essential oils with antibacterial properties, for preventive purpose, to limit health pressure**
- **High level of protein, and high proportion of marine product**
- **Enriched with Vitamins, Minerals, Yeasts and Hepato-protectors to boost the immune system, improve health and reduce stress of freshwater carnivores.**
- **Range supplemented with additives increasing feces stability**
- **Addition of natural pigments, to enhance skin fish color**

	NEO SUPRA			PERCID STARTER	PERCID GROWER	PERCID FINISHER
Commerciale Ref.	M0 / M1 / M2	AL2 / AL3	AL4	1 / 2 / 3	4 / 5	7 / 9
Diameter (mm)	0,15-0,4 / 0,4-0,7 / 0,7-1	0,8 / 1,1	1,4	1,7 / 2,5 / 3,2	4 / 5,5	7,5 / 9,5
Presentation	Crumble	Pellet	Pellet	Pellet	Pellet	Pellet
Live weight (g)	< 0,5 to 2,5 g		2,5 to 5 g	5 à 100 g	100 à 500 g	> 500 g

PACKING

NEO SUPRA : 20 kg bag
PERCID STARTER, PERCID GROWER and PERCID FINISHER : 15 or 20 kg bag, bigbag or bulk

Store feed in a cool and dry place

FLOATING TYPE

NEO SUPRA : sinking slowly
PERCID STARTER, PERCID GROWER and PERCID FINISHER : half-floating and floating

INDICATIVE NUTRITIONAL PROFILE

	NEO SUPRA	PERCID STARTER	PERCID GROWER	PERCID FINISHER
	M0 / M1 / M2 / AL2 / AL3 / AL4	1 / 2 / 3	4 / 5	7 / 9
Protein (%)	58	55	52	50
Fats (%)	13	12	15	17
Gross Energy (MJ/Kg)	21,2	20,4	21,4	21,9
Starch (%)	10,1	9,7	10,3	10
Fibre (%)	0,5	0,9	1,5	1,8
Ash (%)	9	10,5	7,8	7,5
Phosphorus (%)	1,5	1,5	1,2	1,1

VITAMINS INCORPORATION

Vit. A (UI/Kg)	12 000
Vit. D3 (UI/Kg)	2 100
Vit. E (mg/Kg)	440
Vit. C (mg/Kg)	1 000

SUSTAINABILITY

Here at Le Gouessant, we believe that aquaculture is sustainable if performance, quality and preservation of marine resources are combined, that is why we are Global GAP certified.

FEEDING TABLE

Live Weight (g)		Feed	P	F	Diameter		Feeding rate (% of Biomass / day) following water temperature													
From	To				N°	(mm)	18	19	20	21	22	23	24	25	26	27	28	29	30	
<0,1					AL00	< 0,15	<i>Ad libitum</i>													
0,1		0,5	58	13	AL0	0,15-0,4	<i>Ad libitum</i>													
0,5		1,5			AL1	0,4-0,7	3,21	3,49	3,80	4,11	4,41	4,74	5,05	5,44	5,53	4,93	3,90	2,74	1,88	
1,5		2,5			AL2	0,7-1	2,74	2,98	3,27	3,55	3,84	4,08	4,37	4,66	4,79	4,23	3,46	2,66	1,87	
2,5		5			AL3	1,2	2,42	2,62	2,87	3,13	3,39	3,58	3,82	4,00	4,11	3,68	3,11	2,50	1,87	
5		10			AL4	1,4	2,02	2,21	2,39	2,58	2,76	2,94	3,13	3,26	3,33	3,17	2,74	2,29	1,82	
10		15	55	12	1	1,7	1,65	1,74	1,84	1,95	2,05	2,15	2,24	2,31	2,38	2,38	2,32	2,11	1,77	
15		20			2	2,5	1,54	1,64	1,74	1,84	1,93	2,02	2,09	2,15	2,21	2,21	2,21	2,16	1,98	1,72
20		50			2	2,5	1,49	1,59	1,69	1,78	1,88	1,95	2,03	2,07	2,11	2,10	2,10	2,06	1,90	1,67
50		100			3	3,2	1,43	1,54	1,64	1,73	1,82	1,89	1,96	2,00	2,01	1,99	1,99	1,95	1,82	1,61
100		150			3	3,2	1,38	1,48	1,58	1,67	1,76	1,82	1,89	1,92	1,93	1,91	1,87	1,76	1,56	
150		200	52	15	4	4	0,96	1,06	1,17	1,28	1,39	1,50	1,61	1,69	1,76	1,77	1,78	1,64	1,51	
200		300			4	4	0,76	0,85	0,93	1,02	1,11	1,19	1,28	1,34	1,41	1,41	1,41	1,41	1,31	1,20
300		400			5	5,5	0,67	0,74	0,81	0,89	0,97	1,05	1,12	1,18	1,23	1,24	1,24	1,24	1,15	1,05
400		500			5	5,5	0,58	0,65	0,71	0,78	0,85	0,92	0,98	1,03	1,07	1,08	1,08	1,08	1,01	0,93
500		600			5	5,5	0,50	0,55	0,61	0,68	0,74	0,80	0,85	0,89	0,93	0,94	0,94	0,88	0,81	
600		800	50	17	7	7,5	0,44	0,48	0,53	0,58	0,63	0,68	0,73	0,77	0,80	0,81	0,81	0,75	0,69	
800		1 000			7	7,5	0,36	0,40	0,44	0,48	0,52	0,56	0,60	0,63	0,66	0,66	0,66	0,66	0,62	0,56
					9	9,5	0,29	0,33	0,36	0,39	0,43	0,46	0,49	0,52	0,54	0,54	0,54	0,55	0,50	0,46

Feeding rates are indicatives and based on the feed nutritional values. It must be adapted to local conditions and farming goals