

T-CATFISH RANGE

Extruded feed for Catfish



- Floating pellets with a high water stability
- Supplemented fingerling feed to strengthen their immune system
- Use of essential oils with anti-bacterial properties, to preventive purpose, to limite pathological pressure
- Inclusion of marine products in order to increase palatability and optimize feed efficiency
- High digestibility feed range

| | NEO SUPRA | | CATFISH STARTER | T-CATFISH GROWER | T-CATFISH FINISHER |
|------------------|----------------------------|-----------|-----------------|------------------|--------------------|
| Commerciale Ref. | M0 / M1 / M2 | AL2 / AL3 | 1 / 2 | 3 / 4 / 5 | 5 / 7 / 9 |
| Diameter (mm) | 0,15-0,4 / 0,4-0,7 / 0,7-1 | 0,8 / 1,2 | 1,7 / 2,5 | 3,2 / 4 / 5,5 | 5,5 / 7,5 / 9,5 |
| Presentation | Crumble | Pellet | Pellet | Pellet | Pellet |
| Live weight (g) | < 0,5 to 5 g | | 5 to 50 g | 50 to 1 000 g | 1 000 to >1 500 g |

PACKING

NEO SUPRA : 20 kg bag (or 10 kg bucket)
CATFISH STARTER, GROWER and FINISHER : 15 kg bag
 Store feed in a cool and dry place .

FLOATING TYPE

NEO SUPRA: Half-floating (Y)
CATFISH STARTER, GROWER and FINISHER : Floating (Z)

INDICATIVE PROFILE NUTRITIONAL

| | NEO SUPRA | CATFISH STARTER | T-CATFISH GROWER | T-CATFISH FINISHER |
|-----------------------------|-----------|-----------------|------------------|--------------------|
| Protein (%) | 58 | 46 | 42 | 38 |
| Fats (%) | 12 | 10 | 11 | 12 |
| Gross Energy (MJ/Kg) | 21,2 | 19,5 | 19,7 | 19,7 |
| Starch (%) | 11,3 | 15,6 | 16,5 | 21,8 |
| Fibre (%) | 0,5 | 3 | 3,2 | 3,4 |
| Ash (%) | 9 | 9,2 | 8,7 | 7,7 |
| Phosphorus (%) | 1,5 | 1,3 | 1,2 | 1,1 |

VITAMINS INCORPORATION

| | | |
|-----------------|--------|-------|
| Vit. A (UI/Kg) | 12 000 | 8 000 |
| Vit. D3 (UI/Kg) | 2 100 | 1 400 |
| Vit. E (mg/Kg) | 440 | 160 |
| Vit. C (mg/Kg) | 1 000 | 150 |

SUSTAINABILITY

Here at Le Gouessant, we believe that aquaculture is sustainable if performance, quality and preservation of marine resources are combined, that is why we are Global GAP certified.



FEEDING TABLE

| Live Weight (g) | | Feed | P | F | Diameter | | Feeding rate (% of Biomass / day) following water temperature | | | | | | | | | | | | | | | | | | |
|-----------------|------|--------------------|----|----|----------|------------|---|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| From | To | | | | N° | (mm) | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 |
| < 0,5 | | NEO SUPRA | 58 | 13 | AL0 | 0,15 - 0,3 | Ad libitum | | | | | | | | | | | | | | | | | | |
| 0,5 | 2,5 | | | | AL1 | 0,3 - 0,5 | Ad libitum | | | | | | | | | | | | | | | | | | |
| 2,5 | 5 | | | | AL2 | 0,5 - 0,8 | Ad libitum | | | | | | | | | | | | | | | | | | |
| 5 | 10 | T-CATFISH STARTER | 46 | 10 | AL3 | 1,2 | 1,21 | 1,82 | 2,42 | 3,23 | 4,04 | 4,91 | 5,77 | 6,49 | 7,21 | 7,61 | 8,02 | 8,23 | 8,44 | 8,31 | 8,18 | 7,98 | 7,78 | 7,39 | 7,00 |
| 10 | 50 | | | | 1 | 1,7 | 1,09 | 1,64 | 2,18 | 2,91 | 3,64 | 4,42 | 5,19 | 5,84 | 6,49 | 6,85 | 7,22 | 7,41 | 7,59 | 7,48 | 7,37 | 7,18 | 7,00 | 6,65 | 6,30 |
| 50 | 100 | T-CATFISH GROWER | 42 | 11 | 2 | 2,5 | 0,82 | 1,23 | 1,64 | 2,18 | 2,73 | 3,31 | 3,90 | 4,38 | 4,87 | 5,14 | 5,41 | 5,55 | 5,70 | 5,61 | 5,53 | 5,39 | 5,25 | 4,99 | 4,72 |
| 100 | 150 | | | | 3 | 3,2 | 0,64 | 0,95 | 1,27 | 1,70 | 2,12 | 2,58 | 3,03 | 3,41 | 3,79 | 4,00 | 4,21 | 4,32 | 4,43 | 4,36 | 4,30 | 4,19 | 4,08 | 3,88 | 3,67 |
| 150 | 300 | | | | 5 | 5,5 | 0,41 | 0,61 | 0,82 | 1,09 | 1,36 | 1,66 | 1,95 | 2,19 | 2,44 | 2,57 | 2,71 | 2,78 | 2,85 | 2,81 | 2,76 | 2,69 | 2,62 | 2,49 | 2,36 |
| 300 | 500 | | | | 5 | 5,5 | 0,24 | 0,35 | 0,47 | 0,63 | 0,79 | 0,96 | 1,13 | 1,27 | 1,41 | 1,49 | 1,56 | 1,60 | 1,65 | 1,62 | 1,60 | 1,56 | 1,52 | 1,44 | 1,36 |
| 300 | 500 | T-CATFISH FINISHER | 38 | 12 | 5 | 5,5 | 0,35 | 0,53 | 0,71 | 0,94 | 1,17 | 1,43 | 1,68 | 1,88 | 2,09 | 2,22 | 2,33 | 2,39 | 2,46 | 2,42 | 2,38 | 2,33 | 2,26 | 2,15 | 2,04 |
| 500 | 800 | | | | 5 | 5,5 | 0,38 | 0,57 | 0,77 | 1,02 | 1,27 | 1,54 | 1,82 | 2,04 | 2,26 | 2,39 | 2,51 | 2,59 | 2,66 | 2,61 | 2,57 | 2,51 | 2,44 | 2,32 | 2,20 |
| 800 | 1000 | | | | 7 | 7,5 | 0,29 | 0,44 | 0,59 | 0,79 | 0,98 | 1,19 | 1,40 | 1,58 | 1,75 | 1,85 | 1,94 | 2,00 | 2,05 | 2,02 | 1,99 | 1,94 | 1,89 | 1,79 | 1,70 |
| 1000 | 1500 | | | | 7 | 7,5 | 0,26 | 0,38 | 0,51 | 0,68 | 0,85 | 1,04 | 1,22 | 1,37 | 1,52 | 1,61 | 1,69 | 1,73 | 1,78 | 1,75 | 1,73 | 1,69 | 1,64 | 1,56 | 1,47 |
| > 1500 | | | | | 7 | 7,5 | 0,20 | 0,30 | 0,40 | 0,54 | 0,67 | 0,81 | 0,96 | 1,08 | 1,20 | 1,26 | 1,33 | 1,37 | 1,40 | 1,38 | 1,36 | 1,33 | 1,29 | 1,23 | 1,16 |
| | | | | | 7 | 7,5 | 0,16 | 0,25 | 0,33 | 0,44 | 0,55 | 0,67 | 0,78 | 0,88 | 0,98 | 1,03 | 1,09 | 1,12 | 1,15 | 1,13 | 1,11 | 1,08 | 1,06 | 1,00 | 0,95 |

Feeding rates are indicatives and based on the feed nutritional values. It must be adapted to local conditions and farming goals